

Velocity in Lorentz-Violating Fermion Theories

B. Altschul¹

*Department of Physics
Indiana University
Bloomington, IN 47405 USA*

Don Colladay

*New College of Florida
Sarasota, FL 34243 USA*

Abstract

We consider the role of the velocity in Lorentz-violating fermionic quantum theory, especially emphasizing the nonrelativistic regime. Information about the velocity will be important for the kinematical analysis of scattering and other problems. Working within the minimal standard model extension, we derive new expressions for the velocity. We find that generic momentum and spin eigenstates may not have well-defined velocities. We also demonstrate how several different techniques may be used to shed light on different aspects of the problem. A relativistic operator analysis allows us to study the behavior of the Lorentz-violating *Zitterbewegung*. Alternatively, by studying the time evolution of Gaussian wave packets, we find that there are Lorentz-violating modifications to the wave packet spreading and the spin structure of the wave function.

¹baltschu@indiana.edu

1 Introduction

Recent work has stimulated a great deal of interest in the possibility of there existing small Lorentz- and CPT-violating corrections to the standard model. If violations of these fundamental symmetries exist at low and medium energies, they could represent powerful clues as to the nature of Planck scale physics. Within the context of effective field theory, a general Lorentz-violating extension of the standard model has been developed [1, 2], and the stability [3] and renormalizability [4] of this extension have been carefully studied. However, the general standard model extension (SME) is extremely complicated, and even superficially simple questions about its physics may have subtle and even ambiguous answers. For example, the study of the gauge invariance properties of and finite radiative corrections to this Lorentz-violating field theory has proven to be a fruitful source for new theoretical insights [5, 6, 7, 8, 9]. Many other elementary questions about the SME still remain unanswered.

The SME provides a framework within which to analyze the results of experiments testing Lorentz violation. To date, such experimental tests have included studies of matter-antimatter asymmetries for trapped charged particles [10, 11, 12, 13] and bound state systems [14, 15], determinations of muon properties [16, 17], analyses of the behavior of spin-polarized matter [18, 19], frequency standard comparisons [20, 21, 22], measurements of neutral meson oscillations [23, 24, 25], polarization measurements on the light from distant galaxies [26, 27, 28], and others. The analysis of the relevant experimental data requires a good understanding of the behavior of elementary particles in the presence of Lorentz violation. However, there still remain many aspects of Lorentz-violating quantum mechanics about which much more could be known.

In this paper, we shall examine one important aspect of Lorentz-violating fermionic quantum mechanics: the behavior of the velocity. A detailed understanding of the role of the velocity in Lorentz-violating physics is important for analyses of Lorentz-violating scattering processes. The relevant kinematics of such processes may depend sensitively on the character of the velocity. For example, the question of whether the vacuum Cerenkov reaction [29, 30] $e^- \rightarrow e^- \gamma$ can occur is obviously intimately related to whether or not the initial electron's velocity is superluminal. There have been some prior investigations into the properties of the velocity in the presence of Lorentz violation. Some of these analyses have studied the effects of specific forms of Lorentz violation, involving selected terms from the SME [31] or Lorentz violation through “double Special Relativity” [32, 33, 34]. There has also been some analysis [35] of velocities in nonrelativistic Lorentz-violating theories (obtained from their relativistic counterparts through the use of the Foldy-Wouthuysen transformation [36]). However, there has been no comprehensive treatment of the velocity in the context of the Lorentz-violating SME.

This study is a natural outgrowth of earlier work on the effects of Lorentz violation on scattering. Previously, we have looked at the impact of specific Lorentz-violating coefficients on particular scattering processes [37, 38]. Lorentz violation affects particles'

velocities and hence the kinematics of reactions. These effects can be just as important as any change in the dynamics. Explicit relations between the velocity, momentum, and spin are required for these types of calculations. To date, a systematic treatment of these relations is lacking. It is the purpose of this paper to fill this gap and to obtain a more detailed understanding of the role of the velocity in Lorentz-violating quantum theories.

In this paper, we shall concentrate primarily on the nonrelativistic regime. There are several reasons for doing this. First, this regime has thus far received little attention; previous kinematical studies have tended to focus on relativistic speeds and particular systems, such as ultra-high-energy cosmic rays. Second, only nonrelativistic quantum theory possesses a satisfactory probabilistic interpretation, and we shall need to make use of the probabilistic method when we analyze the time development of wave packets. Third, consideration of the nonrelativistic case will help us to distinguish which of the theory's complications are due to the effects of Lorentz violation and which are merely byproducts of the matrix structure of Dirac theory. For example, the velocities we encounter will generally be spin-dependent, but in the relativistic matrix theory, only the helicity component of the spin is a conserved quantity. If we restrict our attention to the nonrelativistic limit, in which all the spin projections are time-independent, we may avoid this difficulty.

Although we shall mostly be interested in particles with nonrelativistic speeds, we shall use a number of inherently relativistic methods in our study of the velocity. The velocity operator in a relativistic fermion theory has a complicated matrix structure, and we shall examine how this structure is modified by the presence of Lorentz-violating parameters. This analysis will allow us to address questions about aspects of the theory that are inherently relativistic, such as *Zitterbewegung*. We shall also use fully relativistic energy-momentum relations to calculate group velocities. Finally, when it is possible for us to display concise exact expressions for the velocity, these expressions will necessarily be relativistic.

In keeping with our nonrelativistic viewpoint, most of our considerations will apply only to the case of massive particles. Moreover, we shall generally consider only one Lorentz-violating interaction at a time. This is a sensible approach if all the Lorentz-violating parameters are small, so that we need only work to leading order in each parameter. In the presence of multiple forms of Lorentz violation, the various Lorentz-violating contributions to the velocity are then simply additive. However, although we shall concentrate on obtaining the leading order corrections to the velocity, we shall, as noted above, also present exact results when possible. An exact treatment offers substantial additional verification that the effects we uncover are indeed meaningful.

We shall introduce the Lorentz-violating coefficients relevant to a purely fermionic theory in Section 2. Then we shall immediately specialize to a theory containing only one specific Lorentz-violating parameter: a timelike axial vector term. Such a term generates many interesting effects, most of which may be analyzed exactly. We shall begin our study of the velocity with an analysis of the propagating modes of the fermion field in

the presence of this term. From the energy-momentum relation for these modes, we can extract a group velocity. We then move on to a general analysis of all the possible Lorentz-violating terms, using the relativistic Dirac algebra (Section 3). We shall determine the *Zitterbewegung*-free contribution to the velocity operator, which governs the bulk motion of the particles. Finally, we turn in Section 4 to an examination of the time evolution of Gaussian wave packets. This wave packet analysis will allow us to resolve several subtle questions about the role of the velocity that will arise in the course of our discussions. Finally, we shall summarize our conclusions in Section 5.

2 Lorentz-Violating Fermion Theory

2.1 Lagrangian Structure

Since our studies of Lorentz violation will focus only on kinematics, we shall work with a theory that describes a single species of noninteracting fermions. Including only the minimal, superficially renormalizable Lorentz-violating terms, the Lagrange density for this theory is

$$\mathcal{L} = \bar{\psi}(i\Gamma^\mu\partial_\mu - M)\psi, \quad (1)$$

where

$$M = m + \not{a} - \not{b}\gamma_5 + \frac{1}{2}H^{\mu\nu}\sigma_{\mu\nu} + im_5\gamma_5, \quad (2)$$

and

$$\Gamma^\mu = \gamma^\mu + c^{\nu\mu}\gamma_\nu - d^{\nu\mu}\gamma_\nu\gamma_5 + e^\mu + if^\mu\gamma_5 + \frac{1}{2}g^{\lambda\nu\mu}\sigma_{\lambda\nu}. \quad (3)$$

These represent the only superficially renormalizable couplings that are possible in a purely fermionic theory. However, some of the couplings are more interesting than others. In particular, m_5 does not violate Lorentz symmetry, and it may be absorbed into the other coefficients by means of a field redefinition [39]. We shall assume that such a redefinition has already been performed and set $m_5 = 0$. Moreover, e , f , and g are inconsistent with the coupling of the fermion field to standard model gauge fields. However, we shall include these three coefficients for the sake of greater generality.

2.2 Propagation Modes in the Presence of b^0

We shall begin our analysis by considering the specific example of a b -type interaction. This form of interaction is particularly illustrative, and, moreover, the results of our careful analysis of this theory will be needed when we consider the more complicated effects of $d^{\nu\mu}$ in the Appendix. The quantization of this particular system has previously been examined in [40], where some properties of the velocity were noted. In general, the spacetime direction of b is arbitrary, but if b is timelike, we may consider it in an observer frame in which its three spatial components b^i vanish: $b^\mu = (b^0, \vec{0})$. Since the

time dimension plays a special role in the canonical quantization of the Dirac field, this is the most natural frame in which to quantize the theory.

In order to quantize the b -modified theory, we must determine the free propagation modes of the fermion field. So we shall solve the free momentum-space Dirac equation, with the effects of the b term included. Because the matrix γ_5 features prominently in the theory, we shall use the Weyl chiral representation for the Dirac matrices:

$$\gamma^0 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \gamma^j = \begin{bmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{bmatrix}, \gamma_5 = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}. \quad (4)$$

We consider a fermion mode with energy E and three-momentum $\vec{p} = p_3 \hat{z}$. By multiplying by γ^0 , the Dirac equation for this mode may be reduced to

$$\begin{bmatrix} E - b^0 + p_3 \sigma^3 & -m \\ -m & E + b^0 - p_3 \sigma^3 \end{bmatrix} u(p) = 0. \quad (5)$$

If the fermion's spin is quantized along the z -axis, with eigenvalues $\pm \frac{s}{2}$, then we may replace $\sigma^3 \rightarrow s$. The eigenvalue condition for E then becomes

$$E^2 = m^2 + (sp_3 - b^0)^2 = m^2 + (s|\vec{p}| - b^0)^2. \quad (6)$$

Note that even though there is no breaking of rotation invariance, the energy depends upon the spin direction, through the helicity s . The explicit electron spinors corresponding to a specified momentum are given in [38]; the corresponding positron spinors are precisely analogous. These solutions may easily be generalized to describe modes with arbitrary three-momentum, so long as the spin is quantized along the direction of the motion, so that s represents the helicity. In the nonrelativistic limit, both the positive- and negative-energy spinors approach their usual limits, provided that b^0 is small compared to the mass.

A number of odd facts follow from the dispersion relation (6). These properties have been previously noted in [3, 31], but we reiterate them here, because a proper understanding of these somewhat counterintuitive results is necessary if one is to possess a complete understanding of the meaning of the velocity in Lorentz-violating physics. Restricting our attention to the $E > 0$ modes, we see that the $\vec{p} = 0$ mode has energy $\sqrt{m^2 + (b^0)^2}$, but that this is not the lowest-possible energy. Indeed, for $s|\vec{p}| = b^0$, the energy is only m . So for $b^0 \neq 0$, a particle with $\vec{p} = \vec{0}$ can release energy by absorbing momentum.

However, such a particle, with vanishing three-momentum, is not actually stationary. Because of the Lorentz-violation, the group velocity for a wave packet centered around $\vec{p} = \vec{0}$ is nonvanishing. In fact, the group velocity for an arbitrary wave packet that is well-localized in momentum space is

$$\vec{v}_g = \vec{\nabla}_{\vec{p}} E = \frac{\vec{p} - sb^0 \hat{p}}{E}. \quad (7)$$

Note that both the numerator and denominator contain contributions that are first order in b^0 . For vanishing $\vec{p}$ and $b^0 < 0$, the velocity points along the direction of the spin. Since $\vec{v}_g$ is a vector and the spin is an axial vector, this result represents a clear physical manifestation of the parity violation arising from γ_5 . Moreover, if the spin is not quantized along the direction of $\vec{p}$, then the wave function will contain a superposition of two different energy eigenstates, with the two energies corresponding to different group velocities. (It is perhaps somewhat unsurprising that there are momentum eigenstates which do not possess a unique group velocity, because the very concepts that we use to define the velocity—the flows of energy and particle density—lose some of their usual meanings when breakdowns of Lorentz symmetry occur.) This also creates the possibility that the wave packet may bifurcate, separating itself into two distinct spin states. However, since the two spin components perpendicular to the momentum are not constants of the motion in the Dirac theory, neither is it clear that such bifurcation will in fact occur. Only a careful analysis of the problem in terms of wave packets can answer this question. We shall perform such an analysis in Section 4.2; for a wave packet with vanishing mean three-momentum, we find that, when b^0 is small compared with the momentum spread of the wave packet, no bifurcation of the wave packet in the plane perpendicular to the initial spin can occur.

3 Operator Methods

Because the group velocity for a wave packet may, under certain circumstances (as in systems displaying anomalous dispersion) become meaningless or even undefined, some further confirmation that (7) represents a real physical velocity is desirable. This confirmation may be found through an analysis of the operator structure of the Dirac theory.

In Dirac theory, the velocity possesses a number of decidedly nonclassical properties. In the presence of Lorentz violation, this fact is even more evident. Indeed, the Lorentz-violating contributions to the velocity may depend upon operators that have no classical analogs. Moreover, unlike ordinary oscillatory *Zitterbewegung* terms, the Lorentz-violating additions to a fermion's velocity will unavoidably affect the bulk motion of that particle.

We shall examine the effects of all the Lorentz-violating interactions included in (1) in this way, beginning with the M terms and, in particular, the specific $M = m - b^0 \gamma^0 \gamma_5$ considered in Section 2.2. After obtaining an operator result analogous to (7) for the b^0 theory, we shall generalize our methods to deal with the remaining M terms and the Γ^μ terms as well.

3.1 Potential (M) Terms

When $M = m - b^0 \gamma^0 \gamma_5$ and $\Gamma^\mu = \gamma^\mu$, the Heisenberg equation of motion for the position operator x_k is

$$\dot{x}_k = i[H, x_k], \quad (8)$$

where

$$H = \alpha_j p_j + \beta m - \gamma_5 b^0 \quad (9)$$

is the single-particle Dirac Hamiltonian. The Dirac matrices are, as usual, $\alpha_j = \gamma^0 \gamma^j$ and $\beta = \gamma^0$. So the time derivative of x_k is simply α_k , just as in the Lorentz-invariant theory, and $\vec{\alpha}$ remains the correct operator to represent the velocity.

We shall therefore consider the equation governing the time evolution of α_k . This is

$$\dot{\alpha}_k = i[H, \alpha_k] = i \left(-2\alpha_k H + 2p_k - 2b^0 \alpha_k \gamma_5 \right). \quad (10)$$

When $b^0 = 0$ this has the exact solution (keeping in mind that $\vec{p}$ and H are constants of the motion)

$$\alpha_k(t) = p_k H^{-1} + \left[\alpha_k(0) - p_k H^{-1} \right] e^{-2iHt}. \quad (11)$$

When $b^0 \neq 0$, finding the solution becomes much harder. However, we can determine the solution to first order in b^0 if we consider only the case in which $\vec{p} = \vec{0}$; this is again permissible because $\vec{p}$ is conserved. (Note that we are not boosting the theory into the $\vec{p} = \vec{0}$ frame, because this would generate a spacelike component for b . Instead, we are merely restricting our attention to the $\vec{p} = \vec{0}$ subspace of the theory.) In this case, the time evolution of α is given by

$$\alpha_k(t) \approx -\frac{b^0}{m} \Sigma_k \beta + \left[\alpha_k(0) + \frac{b^0}{m} \Sigma_k \beta \right] e^{-2iHt}. \quad (12)$$

That (12) is indeed correct to first order in b^0 becomes quite clear when we observe several facts about the $\vec{p} = \vec{0}$ subspace. The spin operator $\vec{\Sigma} = \vec{\alpha} \gamma_5$ is a constant of the motion, and β is $\mathcal{O}[(b^0)^1]$, when these operators are considered within this subspace. Moreover, when $\vec{p} = \vec{0}$, $H = \beta m + \mathcal{O}[(b^0)^1]$, so the time evolution of the b -linear terms is governed entirely by their commutators with β .

The second term on the right-hand side of (12) contains the usual *Zitterbewegung* oscillations with frequency $\omega = 2|H| = 2\sqrt{m^2 + (b^0)^2}$. However, the Lorentz violation may modify the structure of this *Zitterbewegung*. The eigenvalues of the operator $\alpha_k(0) + \frac{b^0}{m} \Sigma_k \beta$ (as well as those of H) essentially determine the length scale over which the *Zitterbewegung* occurs. The eigenvalues of $\alpha_k(0) + \frac{b^0}{m} \Sigma_k \beta$ are $\pm \sqrt{1 + (b^0/m)^2}$, and since these have no $\mathcal{O}(b^0)$ contribution, we see that there is no leading-order modification of the length scale over which the *Zitterbewegung* motion occurs, at least for particles with vanishing $\vec{p}$. We do see indications of an effect—a lengthening of the scale—at higher order. However, this effect would be exactly compensated by the increase in the frequency of the oscillations from $2m$ to $2\sqrt{m^2 + (b^0)^2}$. In any case, since we have not conducted a systematic expansion to $\mathcal{O}[(b^0)^2]$, we cannot make a definitive statement about the higher-order corrections. If there were a correction to the length scale of the *Zitterbewegung*, the effect

would manifest itself, for example, in the Darwin term in the Hamiltonian. The Darwin term is generated by *Zitterbewegung* and is proportional to the *Zitterbewegung* scale squared.

However, the first term on the right-hand side of (12) is more interesting. It has precisely the form that we would expect on the basis of (7). It does not oscillate and is not related to interference between positive and negative energy solutions, so it will not be swamped by the ordinary *Zitterbewegung*, nor can it be eliminated by considering wave packets with only positive-energy components. On the basis of this result, we may conclude that the velocity $\vec{v}_g$ is not merely an artifact of the approximations used in the calculation of the group velocity; it is a real effect deriving from the peculiar relativistic quantum mechanics of this system.

The result (12) also demonstrates that velocity and momentum are fundamentally different objects in this theory. It is already familiar from the ordinary Dirac theory that the velocity operator has a number of unattractive features. α_k and α_l do not commute with one-another for $k \neq l$, nor do they commute with the Hamiltonian, and the only eigenvalues of α_k are ± 1 . However, if we choose to restrict our attention to wave packets containing only positive energy components or to ignore the rapidly oscillating part of $\alpha_k(t)$ (which averages to zero), we can sensibly identify the k -th component of the velocity with the first term in (11), $p_k H^{-1}$, which is exactly the classical velocity for a relativistic particle. However, in (12), there is a new, spin-dependent velocity term that exists even at zero momentum. This term can have no classical analogue, because of its nontrivial matrix structure. There is therefore no way that we can extend the classical correspondence between momentum and velocity to this case. Moreover, it is the unavoidable matrix structure of the velocity that leads to the existence of momentum eigenstates that are not eigenstates of $\vec{V}$.

We may adapt the approximate solution (12) for $\alpha_k(t)$ to account for a much more general M . However, we shall see that, of the possible terms included in M , only a^j , b^0 , and H^{0j} generate nontrivial changes to $\vec{\alpha}$ at leading order. (Beyond leading order, any Lorentz-violating interaction that modifies the energy-momentum relation will cause a corresponding change in the classical velocity term $p_k H^{-1}$.) In order to derive our generalization of (12), let us consider a single-particle Hamiltonian whose $\vec{p} = \vec{0}$ restriction is $H = \beta m + H'$. H' is the source of the Lorentz violation, which is parameterized by some small quantity ϵ : $H' = \mathcal{O}(\epsilon)$. First, let us note that if H' commutes with both α_k and β , then H' is a constant of the motion that does not contribute to $\dot{\alpha}_k$. Of the sixteen possible Dirac matrices, four— I , $\epsilon_{jkl}\sigma^{jl}$, and $\gamma^j\gamma_5$ for $j \neq k$ —have this property. So if H' is any of these, the $\vec{p} = \vec{0}$ solution for α_k is

$$\alpha_k(t) = \alpha_k(0)e^{-2i\beta m t}, \quad (13)$$

and this result is exact.

Otherwise, if either $[H', \beta]$ or $[H', \alpha_k]$ is nonzero, the equation of motion for α_k becomes

$$\dot{\alpha}_k = -2i\alpha_k H + i\{H, \alpha_k\} = -2i\alpha_k H + i\{H', \alpha_k\}. \quad (14)$$

We consider a possible solution to (14), valid through $\mathcal{O}(\epsilon)$ and similar in form to (12):

$$\alpha_k(t) \approx V_k + [\alpha_k(0) - V_k] e^{-2iHt}, \quad (15)$$

where V_k is $\mathcal{O}(\epsilon)$, and $\dot{V}_k$ is $\mathcal{O}(\epsilon^2)$. Inserting this expression into (14), we see that the solution is valid to $\mathcal{O}(\epsilon)$ if $2iV_k H = i\{H', \alpha_k\}$. Therefore, up to corrections of $\mathcal{O}(\epsilon^2)$,

$$V_k \approx \frac{1}{2} \{H', \alpha_k\} H^{-1} \approx \frac{1}{2m} \{H', \alpha_k\} \beta. \quad (16)$$

We must now check the consistency of the statement that $\dot{V}_k = \mathcal{O}(\epsilon^2)$. This amounts to checking that $[\{H', \alpha_k\}, \beta] = 0$. We have already considered all the Dirac matrices that commute with both α_k and β . Therefore, since $[\{H', \alpha_k\}, \beta] = -[\{H', \beta\}, \alpha_k]$, and any two of the basic Dirac matrices either commute or anticommute, all twelve of the remaining matrices— β , γ^j , α_j , σ^{jk} , $\beta\gamma_5$, $\gamma^k\gamma_5$, and γ_5 —satisfy the condition. However, only those of them that commute with α_k will contribute to V_k . So it is precisely those matrices which anticommute with β and commute with α_k that will ultimately affect the velocity.

In fact, in terms of a general $H' = \beta(M - m)$, the leading-order contribution to the velocity is

$$V_k = -\frac{b^0}{m} \Sigma_k \beta - \frac{a^k}{m} \beta + \frac{H^{0j}}{m} \epsilon^{jkl} \Sigma_l. \quad (17)$$

The presence of β in the a and b terms has a simple explanation. To this order, β is simply the sign of the energy, and so it is the operator that determines the direction of the motion.

The three terms in (17) exhaust essentially all the possible contributions to the velocity that can be composed from three-vectors only. At $\vec{p} = \vec{0}$, the only three-vectors that can appear in $\vec{V}$ are the spin, an externally prescribed vector, and a cross product of the spin and a prescribed vector. Each of these types is represented in (17). We also note that, like the b^0 contribution, the a^k and H^{0j} terms in V_k are exactly what could be expected from the energy-momentum relations. In the presence of $\vec{a}$ only, the energy becomes

$$E^2 = m^2 + (\vec{p} - \vec{a})^2, \quad (18)$$

and with only an H^{01} term, the energy is

$$E^2 = m^2 + \vec{p}^2 + (H^{01})^2 \pm 2H^{01} \sqrt{p_2^2 + p_3^2}, \quad (19)$$

where the sign of the square root depends upon the spin orientation. Each of these gives rise to a group velocity in agreement with the corresponding term in (17).

If we repeat our earlier analysis of the *Zitterbewegung* for a^k and H^{0j} , we find that neither of these makes a first-order contribution to the *Zitterbewegung* scale. However, the higher order cancellation we observed does not occur in these cases. The eigenvalues of the

Zitterbewegung component of the velocity depend on the direction under consideration, but the Hamiltonian does not, so the motion could take on an elliptic, rather than spherically symmetric, character. (Again, however, a higher-order calculation would be required in order to verify this.)

We must now discuss the generalization of (17) to the $\vec{p} \neq \vec{0}$ case. There can exist additional complications in the interpretation of the velocity in this situation; for example, as we saw in the theory containing the b^0 term only, there is no unique group velocity unless the spin and the momentum are collinear. However, to the extent that we may approximate $\{H, \alpha_k\}$ to be a constant of the motion, we may also approximate

$$\alpha_k(t) \approx \frac{1}{2} \{H, \alpha_k\} H^{-1} + \left[\alpha_k(0) - \frac{1}{2} \{H, \alpha_k\} H^{-1} \right] e^{-2iHt}. \quad (20)$$

For the b^0 -only theory, (20) is exact for the component of $\vec{\alpha}$ that points parallel to $\vec{p}$, but it need not even be a good approximation for the two perpendicular components.

Fortunately, however, a complete generalization of the operators methods leading to (17) is unnecessary. We may instead utilize observer Lorentz symmetry to determine the velocity when the three-momentum is nonzero. Since we have considered the most general possible M , we may always boost a particle under consideration into a frame in which $\vec{p} = \vec{0}$ (although it would now be a misnomer to refer to this as the particle's "rest frame") and then use (17) to determine $\vec{V}$ in this frame. An inverse boost then gives the velocity in the original frame. This boosting procedure does not require prior knowledge of the particle's velocity, because we are boosting into a frame in which $\vec{p} = \vec{0}$, rather than into one where the particle is stationary. (However, since the velocities involved are $\pm \vec{p}/E$, a knowledge of the Lorentz-violating expression for the energy is required.) This therefore concludes our analysis of the M terms.

3.2 Kinetic (Γ) Terms

We now turn our attention to the Lorentz-violating terms present in Γ^μ . The canonical quantization of the fermion field requires some care when the c , d , e , f , or g coefficients are nonvanishing, because $\mathcal{L}$ may include additional time derivative terms beyond the usual one. A matrix transformation $\psi \rightarrow A\psi$ may be required in order to ensure that $\Gamma^0 = \gamma^0$. (An explicit expression for the required A , valid to all orders in the Lorentz violation, is given in [41].) In this section, we shall assume for simplicity that any such necessary transformation has already been performed. This amounts to setting $c^{\nu 0} = d^{\nu 0} = e^0 = f^0 = g^{\lambda \nu 0} = 0$. However, once such a transformation has been made, we may not freely boost the theory into another frame; such a boost would reintroduce the troublesome contributions to Γ^0 . We will therefore be restricted to considering any theory with a nonstandard Γ^μ in only a single observer frame.

In this section, we shall only consider the terms c , e , and f . The d and g terms, because of their dependences on γ_5 , lead to more complicated calculations; the velocities

will necessarily be spin dependent, and this leads to awkward initial conditions. However, for the theoretically better motivated d term, a complete analysis is given in the Appendix. We shall not consider any *Zitterbewegung* effects in this section, because simple eigenvalue estimates of the distance scales involved do not work when the momentum $\vec{p}$ is nonzero; even in the absence of Lorentz violation, it is very difficult to estimate the scale of the *Zitterbewegung* motion directly for a fermion with nonzero momentum.

In fact, the velocity in the presence of any of c , e , or f may be found exactly. We begin with the fairly straightforward c terms. In the presence of $c^{\nu\mu}$, the single-particle Hamiltonian becomes

$$H = \alpha_j p_j - c_{lj} \alpha_l p_j - c_{0j} p_j + \beta m, \quad (21)$$

so the time derivative of the position operator is

$$\dot{x}_k = \alpha_k - c_{lk} \alpha_l - c_{0k}. \quad (22)$$

The velocity and $\vec{\alpha}$ are no longer one and the same, but the two remain very closely related.

We may solve for α_k with the same methods as we used previously. The equation of motion

$$\dot{\alpha}_k = i [-2\alpha_k (H + c_{0j} p_j) + 2p_k - 2c_{kj} p_j] \quad (23)$$

has the exact solution

$$\alpha_k(t) = (p_k - c_{kj} p_j) (H + c_{0j} p_j)^{-1} + [\alpha_k(0) - (p_k - c_{kj} p_j) (H + c_{0j} p_j)^{-1}] e^{-2i(H + c_{0j} p_j)t}. \quad (24)$$

The *Zitterbewegung*-free contribution to the velocity is therefore

$$V_k = (p_k - c_{kj} p_j - c_{jk} p_j + c_{jk} c_{jl} p_l) (H + c_{0j} p_j)^{-1} - c_{0k}. \quad (25)$$

Note the presence of the inverse Hamiltonian. Whenever H appears in V_k , we must, as we noted following equation (7), account for the Lorentz-violating modifications of the energy eigenvalues when determining the leading-order corrections to the velocity. To first order, only the symmetric part of c_{kj} contributes to $\vec{V}$; the antisymmetric part corresponds at this order merely to a change in the representation of the Dirac matrices, which should have no physical consequences.

In the presence of e^μ or f^μ interactions, with Hamiltonians

$$H = \alpha_j p_j + \beta(m - e_j p_j) \quad (26)$$

or

$$H = \alpha_j p_j + \beta m - i\beta\gamma_5 f_j p_j, \quad (27)$$

the solution (11) for $\alpha_k(t)$ remains unchanged. The only Lorentz-violating contributions to $\vec{V}$ come from the modified relations $\dot{x}_k = \alpha_k - \beta e_k$ and $\dot{x}_k = \alpha_k - i\beta\gamma_5 f_k$. The

Zitterbewegung-free contributions to $\beta(t)$ and $(\beta\gamma_5)(t)$ in the presence of e and f , respectively, are $(m - e_j p_j)H^{-1}$ and $if_j p_j H^{-1}$. So the *Zitterbewegung*-free velocities are

$$V_k = p_k H^{-1} - e_k (m - e_j p_j) H^{-1} \quad (28)$$

and

$$V_k = p_k H^{-1} + f_k (f_j p_j) H^{-1}. \quad (29)$$

Note that the expression (29) for the velocity in the presence of f contains no first-order contributions. These exact results are again entirely in keeping with the energy-momentum relations $E^2 = (m - \vec{e} \cdot \vec{p})^2 + \vec{p}^2$ and $E^2 = m^2 + \vec{p}^2 + (\vec{f} \cdot \vec{p})^2$.

This concludes our analysis of the Dirac algebra relating to the velocity.

4 Wave Packet Analyses

The study of the Dirac algebra associated with the velocity can be quite illuminating, but there remain some questions that can be more satisfactorily answered through the use of other techniques. In particular, we have not dealt with the issue of wave packet bifurcation that was raised earlier. There are also problems relating to wave packet spreading, such as whether ordinary spreading might swamp the Lorentz-violating effects that we are considering. It will also be instructive to examine how the spreading itself may be modified by the Lorentz-violating parameters. To study these aspects of the problem, we must construct and study the particle wave packets directly.

For our wave packet analyses, we shall use the nonrelativistic Foldy-Wouthuysen formulation of the theory [36, 42]. We do this, in part, because only the nonrelativistic quantum theory has a completely consistent probabilistic interpretation. Also, in a nonrelativistic theory, with m large compared to any energy scale associated with the Lorentz-violation, we do not need to include the effects of the negative energy modes; this eliminates the troublesome *Zitterbewegung*, whose structure we have already considered using the operator formalism. Moreover, since the positive-energy spinor $u^s(p)$ is given approximately by

$$u^s(p) \approx \begin{bmatrix} m\chi_s \\ m\chi_s \end{bmatrix}, \quad (30)$$

in the nonrelativistic limit, we shall only need to consider two-component spinor wave functions.

In Section 4.1, we shall derive an expression for the velocity operator in the nonrelativistic Foldy-Wouthuysen representation. We shall then examine how the modified velocity affects the bulk motion and spreading of an initially Gaussian wave packet. However, we shall ignore all spin-dependent effects, except those which contribute directly to the mean velocity. Subsequently, in Section 4.2, we shall treat the problems related to the spin more carefully, examining the possibilities for spin flips and spin-driven wave packet bifurcation.

4.1 Wave Packet Velocities and Spreading

We begin with the nonrelativistic (Foldy-Wouthuysen) single-particle Hamiltonian, as derived in [42].² This is

$$\begin{aligned}
H_{FW} = & \frac{p^2}{2m} + \left[m \left(-c_{jk} - \frac{1}{2} c_{00} \delta_{jk} \right) \right] \frac{p_j p_k}{m^2} \\
& + \left(-b_j + m d_{j0} - \frac{1}{2} m \epsilon_{jkl} g_{kl0} + \frac{1}{2} \epsilon_{jkl} H_{kl} \right) \sigma^j + [a_j - m(c_{0j} + c_{j0}) - m e_j] \frac{p_j}{m} \\
& - \left[b_0 \delta_{jk} - m(d_{kj} + d_{00} \delta_{jk}) - m \epsilon_{klm} \left(\frac{1}{2} g_{mlj} + g_{m00} \delta_{jl} \right) - \epsilon_{jkl} H_{l0} \right] \frac{p_j \sigma^k}{m} \\
& + \left\{ \left[m(d_{0j} + d_{j0}) - \frac{1}{2} \left(b_j + m d_{j0} + \frac{1}{2} m \epsilon_{jmn} g_{mn0} + \frac{1}{2} \epsilon_{jmn} H_{mn} \right) \right] \delta_{kl} \right. \\
& \left. + \frac{1}{2} \left(b_l + \frac{1}{2} m \epsilon_{lmn} g_{mn0} \right) \delta_{jk} - m \epsilon_{jlm} (g_{m0k} + g_{mk0}) \right\} \frac{p_j p_k \sigma^l}{m^2}. \tag{31}
\end{aligned}$$

(Note that [42] uses a different sign convention for p_j and p^j than we do.) Although we neglected them in Section 3, we have included in H_{FW} all possible contributions arising from a nonzero $\Gamma_1^0 = \Gamma^0 - \gamma^0$. It is convenient to combine similar terms in (31) and write the Hamiltonian in the form

$$H_{FW} = (1 - c_{00}) \frac{p^2}{2m} - c_{(jk)} \frac{p_j p_k}{2m} + \frac{\tilde{a}_j}{m} p_j - B_j \sigma^j - B_{jk} p_j \sigma^k + G_{ijk} \frac{p_i p_j \sigma^k}{m}. \tag{32}$$

The notation (jk) represents symmetrization with respect to the indices j and k , defined for example, as $c_{(jk)} = c_{jk} + c_{kj}$. Note that, in any fixed frame, c_{00} and the trace components of c_{jk} may be absorbed into the mass of the fermion and therefore are not observable; we shall henceforth ignore these terms.

The nonrelativistic velocity operator can be computed directly from this Hamiltonian, as

$$\vec{v} = i [H, \vec{x}] = - [H, \vec{\nabla}_{\vec{p}}] = (\vec{\nabla}_{\vec{p}} H). \tag{33}$$

Explicitly, it is given by

$$v_k = \frac{p_k}{m} + c_{(kj)} \frac{p_j}{m} + \frac{\tilde{a}_k}{m} - B_{kj} \sigma^j + G_{(jk)l} \frac{p_j}{m} \sigma^l, \tag{34}$$

where the constants $\tilde{a}_j$, B_{jk} , and G_{jkl} are defined to be

$$\tilde{a}_j = a_j - m(c_{0j} + c_{j0}) - m e_j, \tag{35}$$

$$B_{jk} = \frac{1}{m} \left[b_0 \delta_{jk} - m(d_{kj} + d_{00} \delta_{jk}) - m \epsilon_{klm} \left(\frac{1}{2} g_{mlj} + g_{m00} \delta_{jl} \right) - \epsilon_{jkl} H_{l0} \right], \tag{36}$$

²The same methods could be applied to the fully relativistic Foldy-Wouthuysen Hamiltonian given in [42] as well.

and

$$G_{jkl} = \frac{1}{m} \left\{ \left[m(d_{0j} + d_{j0}) - \frac{1}{2} \left(b_j + md_{j0} + \frac{1}{2} m\epsilon_{jmn} g_{mn0} + \frac{1}{2} \epsilon_{jmn} H_{mn} \right) \right] \delta_{kl} \right. \\ \left. + \left(b_l + \frac{1}{2} m\epsilon_{lmn} g_{mn0} \right) \delta_{jk} - m\epsilon_{jlm} (g_{m0k} + g_{mk0}) \right\}. \quad (37)$$

The result (34) agrees with the nonrelativistic limit of all our other calculations. The same velocity may also be found by taking the direct Foldy-Wouthuysen transform of the relativistic velocity operator.

Now we may determine how the modified velocity $\vec{v}$ affects the bulk motion and spreading of a wave packet. Suppose an initial wave packet is specified as $\psi(\vec{r}, t=0)$. The subsequent time dependence follows from the use of the time evolution operator:

$$\psi(\vec{r}, t) = e^{-iH_{FW}t} \psi(\vec{r}, 0) = e^{-iH_{LV}t} \psi^{(0)}(\vec{r}, t), \quad (38)$$

where H_{LV} is the Lorentz-violating portion of the Hamiltonian and $\psi^{(0)}(\vec{r}, t)$ is the conventional time-dependent wave packet in the absence of Lorentz violation. The decomposition (38) is valid because H_{LV} commutes with the conventional Hamiltonian $\frac{p^2}{2m}$. To lowest order in the violation parameters, the exponential $e^{-iH_{LV}t}$ can be expanded to yield

$$\psi(\vec{r}, t) \approx \psi^{(0)}(\vec{r}, t) - itH_{LV}\psi^{(0)}(\vec{r}, t). \quad (39)$$

As an example, we consider a $\langle \vec{p} \rangle = 0$, spherically symmetric packet with its spin directed along the z -axis ($s = \pm 1$) and an initial spread determined by Δ . The corresponding wave function is

$$\psi^{(0)}(\vec{r}, t) = \left[\frac{\Delta}{\sqrt{\pi} \left(\Delta^2 + i\frac{t}{m} \right)} \right]^{\frac{3}{2}} \exp \left[-\frac{r^2}{2 \left(\Delta^2 + i\frac{t}{m} \right)} \right] \chi_s, \quad (40)$$

where χ_s is the two-component spinor appropriate for the spin state.

We may calculate the action of H_{LV} on the wave packet using the relations

$$p_j \psi^{(0)}(\vec{r}, t) = \left[-i\vec{\nabla}_{\vec{r}} \right]_j \psi^{(0)}(\vec{r}, t) = i \frac{r_j}{\left(\Delta^2 + i\frac{t}{m} \right)} \psi^{(0)}(\vec{r}, t) \quad (41)$$

and

$$p_j p_k \psi^{(0)}(\vec{r}, t) = \frac{1}{\left(\Delta^2 + i\frac{t}{m} \right)} \left[\delta_{jk} - \frac{r_j r_k}{\left(\Delta^2 + i\frac{t}{m} \right)} \right] \psi^{(0)}(\vec{r}, t). \quad (42)$$

This gives a probability density of

$$|\psi(\vec{r}, t)|^2 \approx |N(t)|^2 \exp \left\{ -\frac{\Delta^2}{\left(\Delta^4 + \frac{t^2}{m^2} \right)} \left[(\vec{r} - \vec{v}^s t)^2 + \frac{2t^2 (c_{(jk)} - sG_{(jk)3})}{m^2 \left(\Delta^4 + \frac{t^2}{m^2} \right)} r_j r_k \right] \right\}, \quad (43)$$

where $v_k^s = \tilde{a}_k/m - sB_{k3}$ is the group velocity of the wave packet and $N(t)$ is a normalization factor. The velocity matches the result of (34) for zero momentum, as expected. The c and G terms contribute ellipsoidal deformations in the shape of the wave packet as it spreads; however, in calculating this probability density, we have implicitly summed over the possible spin states, and therefore any spin modulations present in the wave function do not appear in this formula.

4.2 Spin-Dependent Effects

Although $\vec{v}^s$ is s -dependent, the expression (43) for the probability density does not tell us anything about the spin state of the particle. Since we have seen that there can be a complicated interplay between the velocity and the spin in Lorentz-violating fermion theories, it is worthwhile to look at spin effects in more detail. Our examination of wave packet spin structure will not be completely general; in fact, we shall consider a theory in which b^0 is the only Lorentz-violating parameter. However, this single parameter is sufficient to generate many interesting effects. We shall use a specialized approximation method, involving large time position-momentum correlations and the method of stationary phase, to extract information about this theory.

The Hamiltonian governing the nonrelativistic time evolution in the b^0 theory is

$$H_b = \frac{p^2}{2m} - \frac{pb^0}{m}\sigma_{\vec{p}}, \quad (44)$$

where $\sigma_{\vec{p}} = \vec{\sigma} \cdot \hat{p}$ is the Pauli spinor corresponding to the direction of the momentum. This will act on a Gaussian wave packet, with mean momentum $\langle \vec{p} \rangle = \vec{0}$ and the spin oriented along the positive z -axis. The wave function at zero time is given by

$$\psi_b(\vec{r}, t=0) = \left(\frac{1}{\sqrt{\pi}\Delta} \right)^{\frac{3}{2}} e^{-r^2/2\Delta^2} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad (45)$$

$$= (2\sqrt{\pi}\Delta)^{\frac{3}{2}} \int \frac{d^3p}{(2\pi)^3} e^{i\vec{p}\cdot\vec{r}} e^{-p^2\Delta^2/2} \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad (46)$$

The nonrelativistic approximation requires that the length scale Δ satisfy $m\Delta \gg 1$, and we shall also assume that $b^0\Delta \ll 1$, so that the second term in (44) may be treated as a small correction.

In terms of the eigenspinors of $\sigma_{\vec{p}}$, the wave function is

$$\psi_b(\vec{r}, t=0) = (2\sqrt{\pi}\Delta)^{\frac{3}{2}} \int \frac{d^3p}{(2\pi)^3} e^{i\vec{p}\cdot\vec{r}} e^{-p^2\Delta^2/2} \left(\cos \frac{\theta}{2} \begin{bmatrix} \cos \frac{\theta}{2} \\ e^{i\phi} \sin \frac{\theta}{2} \end{bmatrix} + \sin \frac{\theta}{2} \begin{bmatrix} \sin \frac{\theta}{2} \\ -e^{i\phi} \cos \frac{\theta}{2} \end{bmatrix} \right), \quad (47)$$

where θ and ϕ are the spherical coordinates corresponding to the direction of $\vec{p}$. Therefore, the time-evolved wave function is

$$\begin{aligned} \psi_b(\vec{r}, t) &= (2\sqrt{\pi}\Delta)^{\frac{3}{2}} \int \frac{d^3p}{(2\pi)^3} e^{i\vec{p}\cdot\vec{r}} e^{-p^2\Delta^2/2} e^{-ip^2t/2m} \left(e^{i|\vec{p}|b^0t/m} \cos \frac{\theta}{2} \begin{bmatrix} \cos \frac{\theta}{2} \\ e^{i\phi} \sin \frac{\theta}{2} \end{bmatrix} \right. \\ &\quad \left. + e^{-i|\vec{p}|b^0t/m} \sin \frac{\theta}{2} \begin{bmatrix} \sin \frac{\theta}{2} \\ -e^{i\phi} \cos \frac{\theta}{2} \end{bmatrix} \right) \end{aligned} \quad (48)$$

$$= (2\sqrt{\pi}\Delta)^{\frac{3}{2}} \int \frac{d^3p}{(2\pi)^3} e^{i\vec{p}\cdot\vec{r}} e^{-p^2\Delta^2/2} e^{-ip^2t/2m} \begin{bmatrix} \cos \frac{|\vec{p}|b^0t}{m} + i \sin \frac{|\vec{p}|b^0t}{m} \cos \theta \\ ie^{i\phi} \sin \frac{|\vec{p}|b^0t}{m} \sin \theta \end{bmatrix}. \quad (49)$$

The expression (49) is quite complicated. However, for our analysis of the velocity, we are primarily interested in the probability density function $|\psi_b(\vec{r}, t)|^2$, and that only at large times $t \gg m\Delta^2$. At such times, the different momentum modes will have separated themselves in space. We may therefore identify each point in space with a particular value of the momentum [up to a positional uncertainty of $\mathcal{O}(\Delta)$]. However, this identification will not be the same for the upper (spin up) and lower (spin down) components of the spinor wave function [43].

The correct position-momentum correspondence may be found for each component by the method of stationary phase; writing the wave function as

$$\psi_b(\vec{r}, t) = (2\sqrt{\pi}\Delta)^{\frac{3}{2}} \int \frac{d^3p}{(2\pi)^3} \begin{bmatrix} e^{i\Phi_1(\vec{p}, \vec{r}, t)} \\ e^{i\Phi_2(\vec{p}, \vec{r}, t)} \end{bmatrix}, \quad (50)$$

the position may be determined by solving $\text{Re}\{\vec{\nabla}_{\vec{p}}\Phi_j(\vec{p}, \vec{r}, t)\} = 0$. If we neglect those terms that are smaller than $\mathcal{O}(t)$, we find

$$\vec{r} \approx \frac{\vec{p}t}{m} - \begin{cases} \frac{\vec{p}b^0t}{m} \frac{\cos \theta}{\cos^2 \frac{|\vec{p}|b^0t}{m} + \sin^2 \frac{|\vec{p}|b^0t}{m} \cos^2 \theta} & (\text{spin up}) \\ 0 & (\text{spin down}) \end{cases}. \quad (51)$$

If we can invert this expression, to obtain $\vec{p}$ as a function of $\vec{r}$, we may then determine the spatial probability density, using that fact that the density function in momentum space is time-invariant and known. The dominant contribution to $\vec{p}(\vec{r})$ is $\vec{p} \approx m\vec{r}/t$, and we may neglect the corrections to this expression in the argument $|\vec{p}|b^0t/m$ of any trigonometric functions as long as $t \ll m/(b^0)^2$. Making this approximation, we find

$$\vec{p}(\vec{r}) \approx \frac{m}{t} \left(\vec{r} + \frac{b^0t}{m} \frac{\cos \theta}{\cos^2 b^0r + \sin^2 b^0r \cos^2 \theta} \hat{r} \right) \quad (52)$$

for the spin up component. Note that $\vec{p}$ and $\vec{r}$ point in the same direction, so θ is the polar angle for $\vec{r}$ as well as $\vec{p}$.

Our use of the stationary phase method entails three significant approximations. There are obviously small contributions to the wave function coming from regions of phase space

where Φ_j is not stationary, and these we have neglected. We have also neglected any effects from the imaginary part of $\vec{\nabla}_{\vec{p}}\Phi_j$; this means, for example, that the initial uncertainty Δ cannot contribute to this part of the calculation. (Δ will play an important role in our final expression, however.) Finally, we have neglected any terms that do not grow as $\mathcal{O}(t)$ in the large t (or equivalently, far field) limit. This means neglecting any contributions from $\vec{\nabla}_{\vec{p}}\theta$. The gradient with respect to $\vec{p}$ of θ is a time-independent constant, and its coefficient will be a bounded combination of trigonometric functions. This term therefore does not grow linearly with t when $t \gg m\Delta/b^0$, and so does not represent a contribution to the velocity. All three of these approximations are standard elements of the method of stationary phase, although they manifest themselves in slightly unorthodox fashions in this Lorentz-violating problem.

We also need to know the probabilities for the spin of the particle to be either up or down at a given time. These may be found by taking the magnitudes squared of the matrix factors in (49). Then, since each $\vec{p}$ value has a statistical weight of $e^{-p^2\Delta^2}$ in the momentum-space probability density, the position-space density is simply given by

$$|\psi_b(\vec{r}, t)|^2 \propto J e^{-[p(\vec{r})]^2\Delta^2} \left[\frac{\cos^2 \frac{|\vec{p}(\vec{r})|b^0 t}{m} + \sin^2 \frac{|\vec{p}(\vec{r})|b^0 t}{m} \cos^2 \theta}{\sin^2 \frac{|\vec{p}(\vec{r})|b^0 t}{m} \sin^2 \theta} \right], \quad (53)$$

where $J = |\partial(p_1, p_2, p_3)/\partial(x, y, z)|$ is the Jacobian of the transformation from $\vec{p}$ -integration to $\vec{r}$ -integration. We have continued to use a two-component spinor notation; the two matrix components of (53) represent the probability densities for finding the particle at a given position with a particular spin orientation.

The Jacobian is dominated by the $\vec{r}$ -independent term $J \approx (m/t)^3$, but there are also spin-dependent corrections. These correction are obviously zero for the spin down component, but for the spin up portion of the probability density, they could be non-trivial. Sufficiently far from the origin, however, the corrections may be neglected. An examination of the $\mathcal{O}[(b^0)^1]$ correction to the spin up Jacobian will illustrate why. Since all the off-diagonal terms in the spin up Jacobian matrix (the $\partial p_j/\partial r_k$ for $j \neq k$) are necessarily $\mathcal{O}[(b^0)^1]$, and these terms cannot appear singly in any product that makes a nonzero contribution to the determinant, the diagonal approximation,

$$J \approx \frac{\partial p_1}{\partial x} \frac{\partial p_2}{\partial y} \frac{\partial p_3}{\partial z}, \quad (54)$$

is valid up to corrections that are second order in b^0 . Expanding the partial derivatives, the leading order contribution to the spin up J becomes

$$J \approx \left(\frac{m}{t}\right)^3 \left(1 + \frac{b^0 t}{m} \vec{\nabla} \cdot \frac{\cos \theta}{\cos^2 b^0 r + \sin^2 b^0 r \cos^2 \theta} \hat{r}\right). \quad (55)$$

We may treat the trigonometric functions of $b^0 r$ as constants, since their derivatives will

only contribute at $\mathcal{O}[(b^0)^2]$. So to first order, J is

$$J \approx \left(\frac{m}{t}\right)^3 \left(1 + \frac{2b^0 t}{mr} \frac{\cos \theta}{\cos^2 b^0 r + \sin^2 b^0 r \cos^2 \theta}\right). \quad (56)$$

The Lorentz-violating correction appears with a factor of $\frac{b^0 t}{mr} \cos \theta$. At higher order, b^0 will always appear either in connection with this same expression, or in $\frac{(b^0)^2 t}{m}$ (which we have already assumed is small). So for $r \gg \frac{b^0 t}{m} \cos \theta$, we may neglect the b^0 -dependent contributions to J .

Collecting all the necessary elements, we find that our final expression for the probability density is

$$|\psi_b(\vec{r}, t)|^2 \propto \left[\frac{\exp \left\{ -\frac{\Delta^2 m^2}{t^2} \left(r + \frac{b^0 t}{m} \frac{\cos \theta}{\cos^2 b^0 r + \sin^2 b^0 r \cos^2 \theta} \right)^2 \right\} (\cos^2 b^0 r + \sin^2 b^0 r \cos^2 \theta)}{\exp \left(-\frac{\Delta^2 m^2}{t^2} r^2 \right) \sin^2 b^0 r \sin^2 \theta} \right]. \quad (57)$$

Along the z -axis, the density profile is

$$|\psi_b(z, t)|^2 \propto \exp \left[-\frac{\Delta^2 m^2}{t^2} \left(z + \frac{b^0 t}{m} \right)^2 \right] \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \quad (58)$$

Along this direction, the condition we have imposed on r —that $r \gg \frac{b^0 t}{m} \cos \theta$ —is quite strong. For large $|z|$, the exponential tails indicate the presence of an increasingly broad peak that moves with velocity $-b^0/m$; however, we expect that the Gaussian structure of the distribution will be significantly disturbed at smaller values of $|z|$ by the correction coming from J . The wave packet spreading occurs more rapidly than the b -induced drift; however, it would be possible in principle (although not in practice) to measure the latter affect.

In the xy -plane (where $r \gg \frac{b^0 t}{m} \cos \theta$ is automatically satisfied), the probability density has a significantly different form. In terms of $\rho = \sqrt{x^2 + y^2}$, the density is

$$|\psi_b(\rho, t)|^2 \propto \exp \left(-\frac{\Delta^2 m^2}{t^2} \rho^2 \right) \begin{bmatrix} \cos^2 b^0 \rho \\ \sin^2 b^0 \rho \end{bmatrix}. \quad (59)$$

(Note that this represents the joint probability density for x and y , conditional that $z = 0$; it is not a density for ρ itself.) It is in the xy -plane that we might expect to observe a bifurcation of the wave packet. Such bifurcation would be expected to manifest itself as follows: Once the distance $b^0 t/m$ becomes larger than the initial positional uncertainty Δ , the peak in $|\psi_b(\rho, t)|^2$ should move away from $\rho = 0$. The relevant times for this occurrence fall within the range of validity of our approximations, $m\Delta/b^0 \ll t \ll m/(b^0)^2$. However, no such effect is in fact evident. We see only unmodified wave packet spreading, along with a sinusoidal modulation of the spin orientation probabilities.

We see here that the b^0 interaction is capable of inducing spin flips, but these flips only appear at $\mathcal{O}[(b^0)^2]$ and higher. This situation is analogous to the *Zitterbewegung* seen in ordinary Dirac theory. In the Dirac theory, the velocity and the spin are not constants of the motion; they undergo oscillations around their classical mean values. In the nonrelativistic, Lorentz-violating theory, a similar situation exists. If we go beyond leading order, then neither the spin nor the velocity are generally conserved. We have seen the consequences of this in our analysis. The spin flips are obviously related to the nonconservation of σ , and the nonconservation of the velocity also has an evident effect. On space and time scales large enough that certain $\mathcal{O}[(b^0)^2]$ terms become important, the Lorentz-violating correction to $\vec{r}(\vec{p})$ ceases to be a linear function of t ; this signifies that the velocity becomes time dependent.

Although we have seen no evidence for wave packet bifurcation in the plane perpendicular to the motion of the expectation value of the position, such bifurcation definitely can occur along a direction parallel to $\langle \vec{p} \rangle$. An initial wave packet

$$\psi_{p_3}(\vec{r}, 0) = \left(\frac{1}{\sqrt{\pi}\Delta} \right)^{\frac{3}{2}} e^{i\langle p_3 \rangle z} e^{-r^2/2\Delta^2} \begin{bmatrix} \zeta_+ \\ \zeta_- \end{bmatrix}, \quad (60)$$

with $\langle p_3 \rangle$ large compared to both Δ^{-1} and b^0 , will split apart under a time evolution governed by (44). The spin-up component propagates in the z -direction with velocity $\frac{\langle p_3 \rangle - b^0}{m} \hat{z}$, while the spin-down component moves with velocity $\frac{\langle p_3 \rangle + b^0}{m} \hat{z}$. If (in contrast with our earlier example) $b^0 \Delta > 1$, then after a time $t \sim m\Delta/b^0$, the spin states will become well separated in space. This represents a sort of Lorentz-violation-induced Stern-Gerlach phenomenon. However, in the time required for the spin components to separate, the center of mass of the wave packet will move a large distance $V_3 t \sim \langle p_3 \rangle \Delta / b^0$.

5 Summary

In this paper, we have considered the role of the velocity in Lorentz-violating quantum theory. In the presence of Lorentz violation, the velocity becomes a fairly complicated object. In order to deal with this complexity, we have used several different techniques in our analysis. The different methods have also been useful in our efforts to tackle some related subsidiary issues. Although a number of these methods, as well as some of our results, are fully relativistic, we have concentrated especially on the nonrelativistic regime.

In the relativistic regime, new complications, such as potentially superluminal speeds, may arise. In this regard, there is a fundamental difference between the velocities associated with the M terms and those associated with Γ^μ . In the presence of M only, $\vec{\alpha}$ remains the velocity operator, and the *Zitterbewegung*-free portion of $\vec{\alpha}$ has a magnitude that is always less than or equal to one, with equality only at $m = 0$. However, the Γ^μ terms do not necessarily give rise to velocities satisfying $|\vec{V}| \leq 1$, and this fact may pose problems for causality [3]. It seems likely that at these high speeds, some new, possibly

nonlocal physics will come into play, which will ensure that some form of causality is preserved.

We have also seen that there are momentum and spin states that have no well-defined velocity, and because of this, the analysis of scattering experiments can become much more complicated. Cross sections for scattering between generic polarization states may not be well-defined. However, this does not rule out the calculation of certain particular cross sections. For example, in [38] it was shown to be possible to perform all orders evaluations of Compton scattering cross sections, but only if a particular polarization basis, for which the group velocities were time-independent, was used.

When the number of Lorentz-violating coefficients is relatively small, it may be reasonable to solve the modified Dirac equation exactly. We did this for the b^0 theory in Section 2.2. A solution of the Dirac equation automatically entails a determination of the energy-momentum relation, from which the group velocity $\vec{v}_g$ may be determined. This method also involves solving for the exact propagation modes of the field, so it is easy to see precisely which particle and antiparticle states have well-defined velocities.

We also used operator techniques, based on the matrix structure of the Dirac equation, to analyze the velocity. This allowed us to demonstrate several important facts. We showed that the group velocity found in Section 2.2 was not merely a result of anomalous dispersion effects. We also demonstrated that the components of the bulk velocity $\vec{V}$ may be unavoidably noncommuting, so it may not be possible for the Lorentz-violating contribution to $\vec{V}$ to have well-defined projections along more than one axis. Finally, we have derived relativistic operator expressions for $\vec{V}$ to at least first order in a , b , c , e , f , and H (and a similar calculation for d is located in the Appendix).

The operator method also allowed us to address some questions about the *Zitterbewegung* motion in these Lorentz-violating theories. We found that to leading order, none of the Lorentz-violating terms present in M changes the scale of the *Zitterbewegung* oscillations.

Fundamentally, any group velocity is really a property of a wave packet, so in Section 4, we looked explicitly at the effects of Lorentz violation on these packets. We first specialized to the nonrelativistic case, introducing the leading-order Foldy-Wouthuysen Hamiltonian H_{FW} . From H_{FW} , it was a simple matter to obtain an expression for the nonrelativistic velocity operator and to establish its dependence on each of the Lorentz-violating parameters, including g . This result (34) will be useful in all subsequent calculations of nonrelativistic, Lorentz-violating velocities.

We then showed explicitly how this velocity arises, by studying the time evolution of a particular wave packet. Not all of the Lorentz-violating coefficients contribute to the bulk velocity, but even those that do not may affect the structure of the packet. The spreading of the wave packet ceases to be spherically symmetric, because it is deformed by the effects of Lorentz violation.

We also identified some interesting spin effects in our analysis. A free particle that begins in a definite spin state may no longer be in that state at later times. The wave

packet may bifurcate if the initial state is not an eigenstate of the velocity, or the spin orientation probabilities may develop spatial oscillations. These results indicate that there is a great deal of richness in the spin structure of these wave functions.

The results we have derived should have a large number of applications in Lorentz-violating physics. We have provided a very general analysis of the character and role of the velocity in the SME, and so in any situation in which information about the asymptotic propagation states of this theory are required, our expressions should prove useful. We anticipate that this may include analyses of scattering data, decays, and particle oscillations.

Acknowledgments

The authors are grateful to V. A. Kostelecký and H. F. Wolfe for helpful discussions. This work is supported in part by funds provided by the U. S. Department of Energy (D.O.E.) under cooperative research agreement DE-FG02-91ER40661.

Appendix: Velocity Operator in the Presence of d

It is possible to use the relativistic operator methods described in Section 3 to study the velocity even when the kinetic term in the action is spin-dependent (i.e. in the presence of d or g). However, the fact that the spin is not conserved for a free Dirac particle makes these analyses less straightforward than those considered above. We shall outline here the calculation for the $d^{\nu\mu}$ case and display the difficulties associated with it. As in Section 3, we assume that $d^{\nu 0} = 0$.

The presence of γ_5 makes the theory with $d^{\nu\mu}$ much more complicated than the similar $c^{\nu\mu}$ theory. Both the single-particle Hamiltonian

$$H = \alpha_j p_j + d_{lj} \alpha_l p_j \gamma_5 + d_{0j} p_j \gamma_5 + \beta m, \quad (61)$$

and the velocity operator

$$\dot{x}_k = \alpha_k + d_{lk} \alpha_l \gamma_5 + d_{0k} \gamma_5. \quad (62)$$

involve γ_5 . Since they have different Lorentz structures, we shall consider d_{kj} and d_{0j} separately.

In the presence of d_{kj} , the equation of motion for α_k is

$$\dot{\alpha}_k = i (-2\alpha_k H + 2p_k + 2d_{kj} p_j \gamma_5). \quad (63)$$

To first order in d , we may use the Lorentz-invariant expression

$$\gamma_5(t) = \Sigma_j p_j H^{-1} + [\gamma_5(0) - \Sigma_j p_j H^{-1}] e^{-2iHt} \quad (64)$$

for γ_5 in (63). The equation then has the leading order solution

$$\begin{aligned} \alpha_k(t) \approx & p_k H^{-1} + (d_{kj} p_j) (\Sigma_l p_l) H^{-2} + \left[\alpha_k(0) - p_k H^{-1} + (d_{kj} p_j) (\Sigma_l p_l) H^{-2} \right] e^{-2iHt} \\ & + 2i(d_{kj} p_j) \left[\gamma_5(0) - (\Sigma_l p_l) H^{-1} \right] t e^{-2iHt}. \end{aligned} \quad (65)$$

The $d_{lk} \alpha_l \gamma_5 = d_{lk} \Sigma_l$ term in $\dot{x}_k$ also contributes to V_k . The time-independent part of $\Sigma_l(t)$ is $\Sigma_l(0) + \frac{i}{2} \epsilon_{lmn} \alpha_m(0) p_n H^{-1}$ [44]. This operator represents the conserved part of the spin, and the $\alpha_m(0)$ -dependent term vanishes in the nonrelativistic limit. So the *Zitterbewegung*-free velocity is in this case

$$V_k \approx p_k H^{-1} + (d_{kj} p_j) (\Sigma_l p_l) H^{-2} + d_{lk} \left[\Sigma_l(0) + \frac{i}{2} \epsilon_{lmn} \alpha_m(0) p_n H^{-1} \right]. \quad (66)$$

Because the velocity is spin-dependent, the initial conditions enter in an unavoidable fashion. As a result, $\vec{V}$, like $\vec{\alpha}$, possesses a nontrivial matrix structure.

We now consider a d_{0j} term. The inclusion of d_{0j} in the action produces a single-particle Hamiltonian very similar to the H we obtained in the presence of a b^0 interaction. The only difference is that the coefficient of γ_5 in H is now momentum-dependent. However, we may utilize our results from the b^0 case to solve for the velocity in this situation as well. Let us fix k ; for the remainder of this paragraph, this index is not to be summed over. If $\vec{p}$ is oriented in the k -direction, and the spin is also quantized along the k -axis, then all our earlier operator results for b^0 continue to hold. In particular, the *Zitterbewegung*-free part of α_k is approximately $p_k H^{-1} + (d_{0j} p_j) \Sigma_k H^{-1} = p_k H^{-1} (1 + d_{0k} \Sigma_k)$, while the mean value of α_l for $l \neq k$ vanishes. Since the velocity (62) contains an additional, $\vec{\alpha}$ -independent, contribution, the total velocity along the k -direction is

$$V_k \approx p_k H^{-1} (1 + 2d_{0k} \Sigma_k), \quad (67)$$

while in an orthogonal direction $l \neq k$, the velocity is

$$V_l \approx d_{0l} (\Sigma_k p_k) H^{-1}. \quad (68)$$

This agrees with results obtained from the energy-momentum relation,

$$E^2 = m^2 + (sp_k + d_{0j} p_j)^2, \quad (69)$$

where s is again the helicity; from (69) we may obtain the exact group velocities

$$(v_g)_k = \frac{p_k (s + d_{0k})^2}{E} \quad (70)$$

$$(v_g)_l = \frac{s d_{0l} p_k}{E}, \quad (71)$$

in agreement with (67) and (68).

The difficulties with these expressions are twofold. First, there is the unattractive dependence on the conserved spin operator $\Sigma_l(0) + \frac{i}{2}\epsilon_{lmn}\alpha_m(0)p_n H^{-1}$ in (66) and the resulting matrix structure of the velocity. Second, there is the fact that (67) and (68) differ in form. We cannot combine these two equations into a single tensorial expression, because we broke the observer rotation symmetry with our choice of a spin quantization axis. While both these difficulties are surmountable, they do show that the operator method becomes more cumbersome in the presence of spin-dependent Lorentz-violating parameters.

References

- [1] D. Colladay, V. A. Kostelecký, Phys. Rev. D **55**, 6760 (1997).
- [2] D. Colladay, V. A. Kostelecký, Phys. Rev. D **58**, 116002 (1998).
- [3] V. A. Kostelecký, R. Lehnert, Phys. Rev. D **63**, 065008 (2001).
- [4] V. A. Kostelecký, C. D. Lane, A. G. M. Pickering, Phys. Rev. D **65**, 056006 (2002).
- [5] S. Coleman, S. L. Glashow, Phys. Rev. D **59**, 116008 (1999).
- [6] R. Jackiw, V. A. Kostelecký, Phys. Rev. Lett. **82**, 3572 (1999).
- [7] M. Pérez-Victoria, Phys. Rev. Lett. **83**, 2518 (1999).
- [8] V. A. Kostelecký, A. G. M. Pickering, Phys. Rev. Lett. **91**, 031801 (2003).
- [9] B. Altschul, Phys. Rev. D **69**, 125009 (2004).
- [10] R. Bluhm, V. A. Kostelecký, N. Russell, Phys. Rev. Lett. **79**, 1432 (1997).
- [11] R. Bluhm, V. A. Kostelecký, N. Russell, Phys. Rev. D **57**, 3932 (1998).
- [12] G. Gabrielse, A. Khabbaz, D. S. Hall, C. Heimann, H. Kalinowsky, W. Jhe, Phys. Rev. Lett. **82**, 3198 (1999).
- [13] H. Dehmelt, R. Mittleman, R. S. Van Dyck, Jr., P. Schwinberg, Phys. Rev. Lett. **83**, 4694 (1999).
- [14] R. Bluhm, V. A. Kostelecký, N. Russell, Phys. Rev. Lett. **82**, 2254 (1999).
- [15] D. F. Phillips, M. A. Humphrey, E. M. Mattison, R. E. Stoner, R. F. C. Vessot, R. L. Walsworth, Phys. Rev. D **63**, 111101 (R) (2001).
- [16] R. Bluhm, V. A. Kostelecký, C. D. Lane, Phys. Rev. Lett. **84**, 1098 (2000).

- [17] V. W. Hughes, M. G. Perdekamp, D. Kawall, W. Liu, K. Jungmann, G. zu Putlitz, Phys. Rev. Lett. **87**, 111804 (2001).
- [18] R. Bluhm, V. A. Kostelecký, Phys. Rev. Lett. **84**, 1381 (2000).
- [19] B. Heckel, *et al.*, in *Elementary Particles and Gravitation*, edited by B. N. Kursonoglu, *et al.* (Plenum, New York, 1999).
- [20] C. J. Berglund, L. R. Hunter, D. Krause, Jr., E. O. Prigge, M. S. Ronfeldt, S. K. Lamoreaux, Phys. Rev. Lett. **75**, 1879 (1995).
- [21] V. A. Kostelecký, C. D. Lane, Phys. Rev. D **60**, 116010 (1999).
- [22] D. Bear, R. E. Stoner, R. L. Walsworth, V. A. Kostelecký, C. D. Lane, Phys. Rev. Lett. **85**, 5038 (2000).
- [23] V. A. Kostelecký, Phys. Rev. D **61**, 016002 (2000).
- [24] Y. B. Hsiung, Nucl. Phys. Proc. Suppl. **86**, 312 (2000).
- [25] K. Abe *et al.*, Phys. Rev. Lett. **86**, 3228 (2001).
- [26] S. M. Carroll, G. B. Field, R. Jackiw, Phys. Rev. D **41**, 1231 (1990).
- [27] S. M. Carroll, G. B. Field, Phys. Rev. Lett. **79**, 2394 (1997).
- [28] V. A. Kostelecký, M. Mewes, Phys. Rev. Lett. **87**, 251304 (2001).
- [29] R. Lehnert, R. Potting, Phys. Rev. Lett. **93**, 110402 (2004).
- [30] R. Lehnert, R. Potting, hep-ph/0408285.
- [31] D. Colladay, Ph.D. thesis, Department of Physics, Indiana University (1998).
- [32] J. Lukierski, A. Nowicki, Acta Phys. Pol. B **33**, 2537 (2002).
- [33] P. Kosinski, P. Maslanka, Phys. Rev. D **68**, 067702 (2003).
- [34] M. Daszkiewicz, K. Imilkowska, J. Kowalski-Glikman, Phys. Lett. A **323**, 345 (2004).
- [35] H. F. Wolfe, B.A. thesis, Division of Natural Sciences, New College of Florida (2003).
- [36] L. L. Foldy, S. A. Wouthuysen, Phys. Rev. **78**, 29 (1950).
- [37] D. Colladay, V. A. Kostelecký, Phys. Lett. B **511** 209 (2001).
- [38] B. Altschul, Phys. Rev. D **70**, 056005 (2004).

- [39] D. Colladay, P. McDonald, J. Math. Phys. **43** 3554 (2002).
- [40] A. A. Andrianov, P. Giacconi, R. Soldati, JHEP **0202**, 030 (2002).
- [41] R. Lehnert, J. Math. Phys. **45**, 3399 (2004) 3399.
- [42] V. A. Kostelecký and C. D. Lane, J. Math. Phys. **40**, 6245 (1999).
- [43] See D. Bohm, *Quantum Theory* (Prentice-Hall, New York, 1951), p. 204–205, and K. Gottfried, *Quantum Mechanics; Volume I: Fundamentals* (Benjamin, New York, 1966), p. 74–75, for examples of exercises making more conventional use of this position-momentum correlation technique.
- [44] B. Thaller, *The Dirac Equation* (Springer-Verlag, Berlin, 1992), p. 18–22.